

Solving Quadratic Functions: Best Method

Take Square Roots $ax^2 + c = 0$ $(ax + c)^2 = 0$	Completing the Square $ax^2 + bx + c = 0$	Factoring $ax^2 + bx + c = 0$	Quadratic Formula $ax^2 + bx + c = 0$
Key Features: <ul style="list-style-type: none"> • Missing "b" term • Doesn't have to = 0 • Get x^2 or the binomial squared by itself before taking the square root • Don't forget \pm 	Key Features: <ul style="list-style-type: none"> • The "b" term is even • Starts with x^2 or has a GCF • Doesn't need to = 0 • After factoring, set both factors = 0 and SOLVE for x. 	Key Features: <ul style="list-style-type: none"> • Must be in standard form and = 0 • Missing "c" term – GCF factor • After factoring, set both factors = 0 and SOLVE for x. 	Key Features: <ul style="list-style-type: none"> • Must be in standard form and = 0 • Not factorable $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$
Examples: $3x^2 - 7 = 47$ $\frac{2}{3}x^2 - 3 = 7$ $5(x - 4)^2 = 125$	Examples: $x^2 + 6x + 5 = 0$ $x^2 - 2x = 24$ $2x^2 - 8x = 14$	Examples: $3x^2 - 3x - 126 = 0$ $2x^2 - 3x = 0$ $4x^2 - 9 = 0$ $2x^2 - 7x = x^2 - 12$	Examples: $x^2 + 5x + 3 = 0$ $3x^2 - 14x = 5$